



Iron (as Iron (III) Hydroxide Polymaltose) - 50 mg

Canalab

RECover the FERRic levels!!

Iron Deficiency?

Polymaltose Iron 50 mg, improves the production of red blood cells, contributing to the formation of hemoglobin and fighting the symptoms of anemia.

Some of the benefits of RecuFerro are:

Easy to use: RecuFerro comes in ready-to-drink ampoules.

Convenient: RecuFerro can be taken during or after meals, increasing compliance with treatment.

Effective: RecuFerro, unlike products containing Ferrous Sulfate, is composed of Ferric Iron, whose structure protects the bioavailability of iron in the body and its absorption is not affected.

Lower incidence of adverse effects: The high level of absorption of ferric iron decreases the appearance of unpleasant side effects such as diarrhea, constipation, nausea, vomiting, cramps or stained teeth.

Safe: Ferric iron has a lower risk of poisoning in case of overdose.



Delicious Flavor



No metallic taste



No fishy smell

RecuFerro® for the prevention and treatment of iron deficiencies in:
Pregnant and/or breastfeeding women.
Children with anemia or growth retardation due to iron deficiency.
People with malnutrition.



Product Name: Recuferro®

Active ingredient and concentration:
Iron (as Iron (III) Hydroxide Polymaltose) - 50 mg

Pharmaceutical Form: Oral Solution.

Presentation: Ampoule 10 mL, 5 ampoules/blister. Box of 2 blisters.

Pharmacological Action: Iron Deficiency.

Indications: Prevention and supportive treatment of iron deficiency anemia. Use for pregnant, lactating and malnourished people. Children with iron deficiency anemia, stunted growth.

Route of Administration: Oral

Dosage: The dose and duration of treatment depend on the level of iron deficiency. The daily dose can be divided into small doses or one dose. It should be taken during the meal or right after the meal. Adults and children over 12 years: 1 ampoule at a time, 2 times a day. Children under 12 years: 1 ampoule at a time, 1-2 times a day. Children under 2 years: Consult your doctor.

Note: Shake well before use.

Shelf life: 36 months.

Instructions for Use: Separate the blister from the blister. Shake well before use. Unscrew and remove the top of the ampoule. Drink directly from the ampoule. Once opened, it must be completely consumed to avoid contamination of the liquid.

Storage Conditions: Place in a dry and cool place. Protect from direct sunlight, at a temperature below 30°C.

Marketing authorization holder: Canalab S.A. Panamá.

Manufactured in: Hanoi – Vietnam.

12 cm

Code bar:



Canalab

www.canal-lab.com / info@canal-lab.com